

Exercise 9: Pontryagin's Minimum Principle

Prof. Dr. Moritz Diehl, Joel Andersson, Sebastien Gros, Andrea Zanelli, Dimitris Kouzoupis,
Yizhen Wang

Consider the following optimal control problem:

$$\begin{aligned} \min_{x(\cdot), u(\cdot)} \quad & \int_0^T x_1(t)^2 + x_2(t)^2 + u(t)^2 dt \\ \text{s.t.} \quad & \dot{x}_1(t) = (1 - x_2(t)^2) x_1(t) - x_2(t) + u(t), \quad x_1(0) = 0 \\ & \dot{x}_2(t) = x_1(t), \quad x_2(0) = 1 \\ & -1 \leq u(t) \leq 1, \end{aligned} \tag{1}$$

where $T = 10$ as before.

In this exercise, we will apply Pontryagin's maximum principle to this problem and solve it with the *indirect single shooting method*.

1. (a) Introduce the *costate* $\lambda(t)$ and write down the Hamiltonian $H(x(t), \lambda(t), u(t))$ of (1).

$$H(x, \lambda, u) = \lambda_1 ((1 - x_2^2) x_1 - x_2 + u) + \lambda_2 x_1 + x_1^2 + x_2^2 + u^2 \tag{2}$$

- (b) Use Pontryagin's maximum principle to derive an expression for the optimal control $u^*(t)$ as a function of $x(t)$ and $\lambda(t)$. Note: $u(t)$ may only be a piecewise smooth function. Tip: How does u enter in the Hamiltonian?

$$u^* = \max \left(\min \left(-\frac{\lambda_1}{2}, 1 \right), -1 \right) \tag{3}$$

- (c) Derive the *costate equations*, i.e. $\dot{\lambda}(t) = \dots$

$$\begin{aligned} \dot{\lambda}_1 &= -\lambda_1 (1 - x_2^2) - 2x_1 - \lambda_2 \\ \dot{\lambda}_2 &= \lambda_1 x_1 + 2x_2 + \lambda_1 \end{aligned} \tag{4}$$

- (d) Derive the terminal conditions for the costate equations.

$$\lambda_1(T) = 0, \quad \lambda_2(T) = 0. \tag{5}$$

- (e) Augment the original equations with the costate equation to form a two-point boundary-value problem (TPBVP) with four differential equations.

$$\begin{aligned} \dot{x}_1 &= (1 - x_2^2) x_1 - x_2 + u, & x_1(0) &= 0, \\ \dot{x}_2 &= x_1, & x_2(0) &= 1, \\ \dot{\lambda}_1 &= -\lambda_1 (1 - x_2^2) - 2x_1 - \lambda_2, & \lambda_1(T) &= 0, \\ \dot{\lambda}_2 &= \lambda_1 x_1 + 2x_2 + \lambda_1, & \lambda_2(T) &= 0 \end{aligned} \tag{6}$$

- (f) Using the provided template, solve the TPBVP with indirect single shooting. Use $[0, 0]$ as your initial guess for the initial costate. To integrate the system, our best chance is to use a variable stepsize integrator for stiff systems, such as the CVODES integrator from the SUNDIALS suite, available in CasADi. Note that the system is only piecewise smooth, which could potentially cause problems in the integrator, but we will ignore this and hope for the best. The resulting nonlinear system of equations is also challenging to solve, and in CasADi, our best bet is to use IPOPT with a dummy objective function (“minimize 0, subject to $g(x) = 0$ ”).