

# Newton Type Optimization (Unconstrained)

Moritz Diehl

University of Freiburg and University of Leuven

*(some slide material was provided by W. Bangerth and K. Mombaur)*

# Aim of Newton type optimization algorithms

$$\min f(x) \quad (x \in \mathbb{R}^n)$$

- Find a local minimizer  $x^*$  of  $f(x)$ , i.e. a point satisfying

$$\nabla f(x^*) = 0$$

# Derivative based algorithms

- **Fundamental underlying structure of most algorithms:**


- choose start value  $x_0$
- for  $i=1, \dots$ :
  - determine direction of search (descent)  $p$
  - determine step length  $\alpha$
  - new iterate  $x_{i+1} = x_i + \alpha p$
  - check convergence

- Optimization algorithms differ in the choice of  $p$  and  $\alpha$

# Basic algorithm:


## Search direction:

choose descent direction  
( $f$  should be decreased)



## Step length:

solve 1-d minimization approximately,  
satisfy Armijo condition



# Computation of step length


- Dream:

- exact line search:  $\alpha^k = \arg \min_{\alpha} f(x^k + \alpha p^k)$

- In practice:

- **inexact line search:**  $\alpha^k \approx \arg \min_{\alpha} f(x^k + \alpha p^k)$
- ensure sufficient decrease, e.g. Armijo condition

$$f(x_k + \alpha_k p_k) \leq f(x_k) + c_1 \alpha_k \nabla f_k^T p_k$$



# How to compute search direction?

- We discuss three algorithms:
  - Steepest descent method
  - Newton's method
  - Newton type methods

# Algorithm 1: Steepest descent method

- Based on first order Taylor series approximation of objective function

$$f(x_k + p_k) = f(x_k) + \underbrace{\nabla f(x_k)^T p_k}_{\text{direction of descent}} + \dots$$

- maximum descent, if



$$\begin{aligned} \frac{\nabla f(x_k)^T p_k}{\|p_k\|} &\rightarrow \text{min!} \\ \Rightarrow p_k &= -\nabla f(x_k) \end{aligned}$$

# Steepest descent method

Choose steepest descent search direction, perform (exact) line search:

$$p^k = -\nabla f(x^k) \quad x^{k+1} = x^k - \alpha^k \nabla f(x^k)$$

search direction is perpendicular to level sets of  $f(x)$





# Convergence of steepest descent method



steepest descent method has *linear convergence*

$$\text{i.e. } \|x^k - x^*\| \leq C \|x^{k-1} - x^*\|$$


- gain is a fixed factor  $C < 1$
- convergence can be very slow if  $C$  close to 1

If  $f(x) = x^T A x$ ,  $A$  positive definite,  $\lambda$  eigenvalues of  $A$ , one can show that

$$\Rightarrow C \approx \frac{\lambda_{\max} - \lambda_{\min}}{\lambda_{\max} + \lambda_{\min}}$$






# Example - steepest descent method




$$f(x, y) = \sqrt[4]{(x - y^2)^2 + \frac{1}{100}} + \frac{1}{100} y^2$$

banana valley function,  
global minimum at  $x=y=0$



# Example - steepest descent method

$$\|x^k - x^*\|$$



*Convergence of steepest descent method:*

- needs almost 35.000 iterations to come closer than 0.1 to the solution
- mean value of convergence constant  $C$ : 0.99995
- at  $(x=4, y=2)$ , there holds

$$\lambda_1 = 0.1, \lambda_2 = 268 \quad \Rightarrow \quad C \approx \frac{268 - 0.1}{268 + 0.1} \approx 0.9993$$

# Algorithm 2: Newton's Method

- Based on **second order Taylor series approximation** of  $f(x)$

$$f(x_k + p_k) = f(x_k) + \underbrace{\nabla f(x_k)^T p_k + \frac{1}{2} p_k^T \nabla^2 f(x_k) p_k}_{\nabla f(x_k)^T p_k + \frac{1}{2} p_k^T \nabla^2 f(x_k) p_k \rightarrow \min!} + \dots$$



$$\Leftrightarrow \nabla^2 f(x_k) p_k = -\nabla f(x_k)$$

„Newton-Direction“  $p_k = -(\nabla^2 f(x_k))^{-1} \nabla f(x_k)$


# Visualization of Newton's method

$p_k$  minimizes quadratic approximation of the objective

$$Q(p^k) = f(x^k) + \nabla f(x^k) p^k + \frac{1}{2} p^{kT} \nabla^2 f(x^k) p^k$$



if quadratic model is good, then take full step with  $\alpha^k = 1$



# Convergence of Newton's method


Newton's method has *quadratic convergence*

$$\text{i.e.} \quad \|x^k - x^*\| \leq C \|x^{k-1} - x^*\|^2$$

This is ***very fast*** close to a solution:




Correct digits double in each iteration!

# Example - Newton's method




$$f(x, y) = \sqrt[4]{(x - y^2)^2 + \frac{1}{100}} + \frac{1}{100} y^2$$

banana valley function,  
global minimum at  $x=y=0$



# Example - Newton's method

$$\|x^k - x^*\|$$




*Convergence of Newton's method:*

- less than 25 iterations for an accuracy of better than  $10^{-7}$ !
- convergence roughly *linear* for first 15-20 iterations since step length  $\alpha_k \neq 1$
- convergence roughly *quadratic* for last iterations with step length

$$\alpha_k = 1$$



# Comparison of steepest descent and Newton



For banana valley example:

- Newton's method **much faster** than steepest descent method (factor 1000)
- Newton's method superior due to higher order of convergence
- steepest descent method converges too slowly for practical applications

# Algorithm 3: Newton type methods

In practice, evaluation of second derivatives  
for the hessian can be difficult!

- approximate hessian matrix  $\nabla^2 f(x^k)$
- often methods ensure that the approximation  $B_k$  is positive definite

$$x^{k+1} = x^k - B_k^{-1} \nabla f(x^k)$$

$$B_k \approx \nabla^2 f(x^k)$$

methods are collectively known as *Newton type methods*

# Newton type variants

Notation:

$$p_k := x_{k+1} - x_k = -B_k^{-1} \nabla f(x^k)$$

- **Steepest Descent:**

$$B_k = I$$

Convergence rate: linear

- **Newton Method:**

$$B_k = \nabla^2 f(x^k)$$

Convergence rate: quadratic

# Newton type variants (continued)

- **BFGS** update (Broyden, Fletcher, Goldfarb, Shanno)

$$B_{k+1} = \underset{B_{k+1}}{\operatorname{argmin}} \|B_{k+1}^{-1} - B_k^{-1}\|_{W,F}^2$$
$$\text{s.t. } B_{k+1} p_k = \nabla f(x^{k+1}) - \nabla f(x^k)$$
$$B_{k+1} = B_{k+1}^T$$

Convergence rate: super-linear

- For Least-Squares Problems: **Gauss-Newton Method**

$$f(x) = \frac{1}{2} \|F(x)\|^2 \quad J(x) = \frac{\partial F(x)^T}{\partial x}$$
$$B_k = J(x^k)^T J(x^k)$$

Convergence rate: linear

# Summary: Unconstrained Newton Type Optimization

- Aim: find **local minima** of smooth nonlinear problems:  $\nabla f(x^*)=0$
- Derivative based methods iterate  $x_{i+1} = x_i + \alpha_i p_i$  with
  - **search direction**  $p_i$  and **step length**  $\alpha_i$  .
  - start at **initial guess**  $x_0$  ,
- Four Newton type methods:
  - steepest descent: intuitive, but slow linear convergence
  - exact Newton's method: very fast quadratic convergence
  - BFGS: fast superlinear convergence
  - Gauss-Newton (only for least-squares): fast linear convergence

# Literature

- **J. Nocedal, S. Wright: Numerical Optimization, Springer, 1999/2006**
- P. E. Gill, W. Murray, M. H. Wright: Practical Optimization, Academic Press, 1981
- R. Fletcher, Practical Methods of Optimization, Wiley, 1987
- D. E. Luenberger: Linear and Nonlinear Programming, Addison Wesley, 1984