Modelling and System Identification – Microexam 1

Prof. Dr. Moritz Diehl, IMTEK, Universität Freiburg November 18, 2014, 8:15-9:15, Freiburg

Surname:		Name:	Matriculation number	:	
Study	:	Studiengang: Ba	achelor Master	laster	
	•	·	e right answer of each question be		
1.	What is the probability density σ^2 ? The answer is $p_X(x) =$			ariable X with mean μ and variance	
	(a) $e^{\frac{(x-\mu)^2}{2\sigma}}$	(b) $e^{-\frac{(x-\mu)^2}{2\sigma}}$		$(d) \qquad e^{-\frac{(x-\mu)^2}{2\sigma^2}}$	
2.	What is the PDF of a variable Z with uniform distribution on the interval $[c,d]$? For $x \in [c,d]$ it has the value:				
	(a) $p_Z(x) = (d-c)$	(b) $p_Z(x) = (c - \epsilon)$	(c) $p_Z(x) = \frac{x}{\sqrt{d-c}}$	$(\mathbf{d}) \qquad p_Z(x) = \frac{1}{d-c}$	
3.	What is the PDF of an n -dimensional normally distributed variable Z with zero mean and covariance matrix $\Sigma \succ 0$? The answer is $p_Z(x) = \frac{1}{\sqrt{(2\pi)^n \det(\Sigma)}} \dots$				
	(a) $e^{-\frac{1}{2}x^T\Sigma x}$	(b) $e^{-\frac{1}{2}x^T\Sigma^{-1}x}$	$(c) e^{\frac{1}{2}x^T \Sigma x}$		
4.			nd covariance matrix $\Sigma \in \mathbb{R}^{n \times n}$. AX . What is the mean μ_Y of Y ?	For a fixed $a \in \mathbb{R}^m$ and $A \in \mathbb{R}^{m \times n}$	
	(a) $Y - a + AX$	(b) $a + Ad$		$(\mathbf{d}) \square a^T A d + d^T \Sigma d$	
5.	Above in Question $\ref{eq:question}$, what is the covariance matrix of Y ?				
	(a) $\int d^T \Sigma d$	(b) $\triangle A\Sigma A^T$	(c) $A^T \Sigma^{-1} A$	$(\mathbf{d}) \square A\Sigma^{-1}A^T$	
6.	Above in Question $\ref{eq:cov}$, which statement is true? $Cov(Y) =$				
	(a) $Y^\top Y - \mu_Y^\top \mu_Y$			$(b) \qquad \mathbb{E}\{YY^{\top}\} - \mu_Y \mu_Y^{\top}$	
	$(c) \square YY^{\top} - \mu_Y \mu_Y^{\top}$			$(d) \square \mathbb{E}\{Y^\top Y\} - \mu_Y^\top \mu_Y$	
7.	(*) Above in Question ??, what is the mean of the matrix valued random variable $Z = YY^T$?				
	(a) $(a+Ad)(a+Ad)^T$			(b) $\Box aa^T + Add^TA^T + A\Sigma A^T$	
	(c) $(a + Ad)(a + Ad)^T + A\Sigma A^T$			$\boxed{ (d) \square aa^T + Add^T A^T }$	
8.	A scalar random variable has the standard deviation y. What is its variance?				
	(a) \sqrt{y}	(b) y ²	(c) _ y	$(d) \qquad y^{-1}$	
9.	A scalar random variable has the variance w . What is its standard deviation?				
	(a) w	(b) w^{-1}	$(c) \square w^2$	(d) \sqrt{w}	
10.	Regard a random variable $\beta \in \mathbb{R}$ with zero mean and variance σ^2 . What is the mean of the random variable $z = \beta^2$?				
	(a) $\beta + \sigma^2$	(b) _ σ	(c)	(d) $\beta + \sigma$	
11.	(*) Regard a random variable $X \in \mathbb{R}^n$ with zero mean and covariance matrix Σ . What is the mean of $Z = X^T X$?				
	(a) $\ \Sigma\ _F^2$	(b) \subseteq $\det(\Sigma)$	$\ (c) \square \ \Sigma \ _2^2$	(d) \square trace(Σ)	

points on page: 11

1

12.	What is the minimizer x^* of the convex function $f:\mathbb{R}\to\mathbb{R}$,	$f(x) = e^x - 2x ?$			
	(a) $x^* = -1$ (b) $x^* = 1$	(c) $x^* = \log_e(2)$ (d) $x^* = 0$			
13.	What is the minimizer x^* of the convex function $f:\mathbb{R}\to\mathbb{R}$,	$f(x) = \alpha + \beta x + \frac{1}{2}\gamma x^2 \text{ with } \gamma > 0 ?$			
	(a) $x^* = \frac{2\beta}{\alpha}$ (b) $x^* = -\frac{\beta}{\gamma}$	(c) $x^* = -\frac{\beta}{2\gamma}$ (d) $x^* = -\frac{\beta}{\alpha}$			
14.	What is the minimizer of the convex function $f: \mathbb{R}^n \to \mathbb{R}$,	$f(x) = \ y - \Phi x\ _2^2$ (with Φ of rank n)? The answer is $x^* = \dots$			
		$(c) \square (\Phi^T \Phi)^{-1} \Phi^T y \qquad (d) \square (\Phi \Phi^T)^{-1} \Phi^T y$			
15.	6. What is the minimizer of the function $f: \mathbb{R}^n \to \mathbb{R}$, $f(x) = \ b + B^T x\ _2^2$ (with B^T of rank n)? The answer is $x^* = a^T x^2 + a^$				
	(a) \square $(BB^T)^{-1}B^Tb$ (b) \square $-(BB^T)^{-1}Bb$	(c) \square $(B^TB)^{-1}B^Tb$ $ \qquad \qquad (d) \qquad \square \qquad -(B^TB)^{-1}B^Tb $			
16.	For a matrix $\Phi \in \mathbb{R}^{N \times d}$ with rank d (and $N \geq d$), what is its	pseudo-inverse Φ^+ ?			
	(a) $ (\Phi \Phi^T)^{-1} \Phi^T $ (b) $ (\Phi \Phi^T)^{-1} \Phi $	$(c) \qquad (\Phi^T \Phi)^{-1} \Phi^T \qquad (d) \qquad (\Phi^T \Phi)^{-1} \Phi$			
17.	7. Given a sequence of numbers $y(1), \dots, y(N)$, what is the minimizer θ^* of the function $f(\theta) = \sum_{k=1}^{N} (y(k) - \theta)^2$?				
	(a)				
18.	What does "i.i.d." stand for?				
	(a) infinite identically disturbed	(b) infinite identically dependent			
	(c) independent identically disturbed	(d) independent identically distributed			
19.	Given a sequence of i.i.d. scalar random variables $X(1), \ldots, X(N)$, each with mean μ and variance σ^2 , what is the exp value of their arithmetic mean, i.e. of the random variable Y defined by $Y = \frac{1}{N} \sum_{k=1}^{N} X(k)$?				
	(a) μ (b) $\frac{\mu}{N}$	$(c) \qquad \qquad$			
20.	In Question $\ref{eq:question}$, what is the variance of the variable Y ?				
	(a) $\frac{\sigma}{N}$ (b) $\frac{\sigma}{N-1}$	$(c) \qquad \qquad \boxed{ (d) \qquad \qquad \frac{\sigma^2}{N}}$			
21.	1. Given a prediction model $y(k) = \theta_1 + \theta_2 x(k)^2 + \epsilon(k)$ with unknown parameter vector $\theta = (\theta_1, \theta_2)^T$, and assuming i.i.d. noise $\epsilon(k)$ with zero mean, and given a sequence of N scalar input and output measurements $x(1), \ldots, x(N)$ and $y(1), \ldots, y(N)$ we want to compute the linear least squares (LLS) estimate $\hat{\theta}_N$ by minimizing the function $f(\theta) = \ y_N - \Phi_N \theta\ _2^2$. If $y_N = (y(1), \ldots, y(N))^T$, how do we need to choose the matrix $\Phi_N \in \mathbb{R}^{N \times 2}$?				
		(c) $\begin{bmatrix} 1 & x(1) \\ \vdots & \vdots \\ 1 & x(N) \end{bmatrix}$ (d) $\begin{bmatrix} 1 & -x(1) \\ \vdots & \vdots \\ 1 & -x(N) \end{bmatrix}$			
		points on page: 10			