

# Modeling and System Identification – Microexam 1

Prof. Dr. Moritz Diehl, IMTEK, Universität Freiburg

November 21, 2018, 14:15-15:15, Freiburg

Surname:

Name:

Matriculation number:

Study:

Programm: Bachelor  Master

Please fill in your name above and tick exactly **ONE** box for the right answer of each question below.  
You can get a maximum of **10 points** on this microexam.

1. Which of the following functions is NOT convex on  $x \in [-1, 1]$

(a) <input type="checkbox"/> $x + 42$	(b) <input type="checkbox"/> $\exp(-x)$	(c) <input type="checkbox"/> $\sin^{-1}(x)$	(d) <input type="checkbox"/> $-\cos(x)$
---------------------------------------	---	---	---

2. What is the probability density function (PDF)  $p_X(x)$  for a normally distributed random variable  $X$  with mean  $-6$  and standard deviation  $3$ ? The answer is  $p_X(x) = \frac{1}{\sqrt{18\pi}} \dots$

(a) <input type="checkbox"/> $e^{-\frac{(x+6)^2}{3}}$	(b) <input type="checkbox"/> $e^{-\frac{(x-6)^2}{9}}$	(c) <input type="checkbox"/> $e^{-\frac{(x-6)^2}{18}}$	(d) <input type="checkbox"/> $e^{-\frac{(x+6)^2}{18}}$
---	---	--	--

3. Which of the following statements does NOT hold for all PDFs  $p(x)$  of a scalar random variable?

(a) <input type="checkbox"/> $\int_{-\infty}^{\infty} p(x)dx = 1$	(b) <input type="checkbox"/> $p(x) \geq 0$
(c) <input type="checkbox"/> $p(x) < 1$	(d) <input type="checkbox"/> $\int_{-1}^1 p(x)dx \geq 0$

4. What is the PDF of a random variable  $Y$  with uniform distribution on the interval  $[0, \sqrt{3}]$ ? For  $z \in [0, \sqrt{3}]$  it has the value:

(a) <input type="checkbox"/> $p_Z(y) = \frac{\sqrt{3}}{y}$	(b) <input type="checkbox"/> $p_Z(y) = \frac{1}{\sqrt{3}}$	(c) <input type="checkbox"/> $p_Y(z) = \frac{1}{\sqrt{3}}$	(d) <input type="checkbox"/> $p_Y(z) = 1$
--	--	--	---

5. Consider a multi-dimensional random variable  $X \in \mathbb{R}^n$  with mean value  $\mu$ . What is the covariance?  $\text{cov}(X) = \dots$

(a) <input type="checkbox"/> $\mathbb{E}\{(X - \mu)\}^2$	(b) <input type="checkbox"/> $\mathbb{E}\{(X - \mu)^2\}$
(c) <input type="checkbox"/> $\mathbb{E}\{(X - \mu)(X - \mu)^\top\}$	(d) <input type="checkbox"/> $\mathbb{E}\{(X - \mu)^\top(X - \mu)\}$

6. Consider a multi-dimensional random variable  $X \in \mathbb{R}^d$ . What are the dimensions of the covariance?  $\text{cov}(X) \in \dots$

(a) <input type="checkbox"/> $\mathbb{R}^{d \times d}$	(b) <input type="checkbox"/> $\mathbb{R}^{d \times d^2}$	(c) <input type="checkbox"/> $\mathbb{R}^{d^2 \times d}$	(d) <input type="checkbox"/> $\mathbb{R}^{d \times 1}$
--	--	--	--

7. Regard a random variable  $X \in \mathbb{R}^n$  with mean  $\mu \in \mathbb{R}^n$  and covariance matrix  $\Sigma \in \mathbb{R}^{n \times n}$ . For a fixed  $b \in \mathbb{R}^n$  and  $D, A \in \mathbb{R}^{m \times n}$ , regard another random variable  $Y \in \mathbb{R}^m$  defined by  $Y = AX + Db$ . The mean of  $Y$  is given by  $\mu_Y = \dots$

(a) <input type="checkbox"/> $A\Sigma^{-1}A^\top$	(b) <input type="checkbox"/> $A^\top Db$	(c) <input type="checkbox"/> $A\mu + Db$	(d) <input type="checkbox"/> $A\mu$
---	--	--	-------------------------------------

8. In the Question above, what is the covariance matrix of  $Y$ ?

(a) <input type="checkbox"/> $D\Sigma^{-1}D^\top$	(b) <input type="checkbox"/> $A^\top \Sigma A$	(c) <input type="checkbox"/> $D^\top \Sigma$	(d) <input type="checkbox"/> $A\Sigma A^\top$
---	--	--	---

9. Regard a random variable  $X \in \mathbb{R}^n$  with zero mean and covariance matrix  $\Sigma \in \mathbb{R}^{n \times n}$ . Given a vector  $c \in \mathbb{R}^n$ , what is the mean of  $Z = c^\top X X^\top c$ ?

(a) <input type="checkbox"/> $\det(\Sigma)$	(b) <input type="checkbox"/> $c^\top \text{trace}(\Sigma) c$	(c) <input type="checkbox"/> $c^\top \Sigma c$	(d) <input type="checkbox"/> $c^\top c \text{trace}(\Sigma)$
---	--	--	--

10. Regard a random variable  $\lambda \in \mathbb{R}$  with zero mean and standard deviation  $d$ . What is the mean of the random variable  $Y = \lambda^2$ ?

(a) <input type="checkbox"/> $0$	(b) <input type="checkbox"/> $d$	(c) <input type="checkbox"/> $d^2$	(d) <input type="checkbox"/> $2\lambda d$
----------------------------------	----------------------------------	------------------------------------	---

11. Regard another scalar random variable that has variance  $(d^2 - 2)$ . What is its standard deviation?

(a) <input type="checkbox"/> 0	(b) <input type="checkbox"/> $d$	(c) <input type="checkbox"/> $\sqrt{d^2 - 2}$	(d) <input type="checkbox"/> $(d^2 - 2)^2$
--------------------------------	----------------------------------	---	--

12. Regard the function  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ ,  $f(x) = \frac{1}{2} \| -b + Dx \|_W^2$  (with  $D$  of rank  $n$  and  $W$  positive definite). What is the gradient of  $f(x)$ ?

(a) <input type="checkbox"/> $Wb$	(b) <input type="checkbox"/> $W(Dx - b)$
(c) <input type="checkbox"/> $(D^T W D)x - D^T W b$	(d) <input type="checkbox"/> $\frac{1}{2} x^T (D^T W D)x - b^T W D x$

13. Regard the function  $f$  from the previous question and the optimization problem  $x^* = \arg \min f(x)$ . The solution is  $x^* =$

(a) <input type="checkbox"/> $(D W D^T)^{-1} D W b$	(b) <input type="checkbox"/> $(D D^T)^{-1} D W b$
(c) <input type="checkbox"/> $(D^T D)^{-1} D^T b$	(d) <input type="checkbox"/> $(D^T W D)^{-1} D^T W b$

14. What is the minimizer  $x^*$  of  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ ,  $f(x) = \frac{1}{2} \| Ax - b \|_2^2$  if  $\text{rank}(A) = n$ ? The solution is  $x^* = \dots$

(a) <input type="checkbox"/> $A^+$	(b) <input type="checkbox"/> $(A^T A)^{-1} A^T b$
(c) <input type="checkbox"/> $A^+ x$	(d) <input type="checkbox"/> $(A^T A)^{-1} A b$

15. Given a sequence of numbers  $y(1), \dots, y(N)$ , what is the minimizer  $\theta^*$  of the function  $f(\theta) = \sum_{k=1}^N (y(k) - 2\theta)^2$ ? The answer is  $\theta^* = \dots$

(a) <input type="checkbox"/> $\frac{1}{2N} \sum_{k=1}^N y(k)^2$	(b) <input type="checkbox"/> $\frac{\sum_{k=1}^N y(k)}{2N}$	(c) <input type="checkbox"/> $\frac{1}{4N} \sum_{k=1}^N y(k)^2$	(d) <input type="checkbox"/> $\sum_{k=1}^N \frac{y(k)}{4N}$
---	---	---	---

16. Consider the model  $y(k) = 2\theta_1 + \frac{\theta_2}{3} x(k)^2 + \frac{\theta_3}{4} x(k)^3 + \epsilon(k)$  and the vector of unknown parameters  $\theta = (\theta_1, \theta_2, \theta_3)^T$ . The additive noise  $\epsilon(k)$  is assumed to have zero mean and to be i.i.d. For a given sequence of  $N$  scalar input and output measurements  $x(1), \dots, x(N)$  and  $y(1), \dots, y(N)$ , we want to compute the linear least squares (LLS) estimate  $\hat{\theta}_N$  by minimizing the function  $f(\theta) = \| y_N - \Phi_N \theta \|_2^2$ . If  $y_N = (y(1), \dots, y(N))^T$ , how do we need to choose the matrix  $\Phi_N \in \mathbb{R}^{N \times 3}$ ?  $\Phi_N = \dots$

(a) <input type="checkbox"/> $\begin{bmatrix} 2 & 3x(1)^2 & 4x(1)^3 \\ \vdots & \vdots & \vdots \\ 2 & 3x(N)^2 & 4x(N)^3 \end{bmatrix}$	(b) <input type="checkbox"/> $\begin{bmatrix} 2 & \frac{x(1)^2}{3} & \frac{x(1)^3}{4} \\ \vdots & \vdots & \vdots \\ 2 & \frac{x(N)^2}{3} & \frac{x(N)^3}{4} \end{bmatrix}$	(c) <input type="checkbox"/> $\begin{bmatrix} \frac{x(1)^2}{3} & 2 & \frac{x(1)^3}{4} \\ \vdots & \vdots & \vdots \\ \frac{x(N)^2}{3} & 2 & \frac{x(N)^3}{4} \end{bmatrix}$	(d) <input type="checkbox"/> $\begin{bmatrix} 2 & x(1)^2 & x(1)^3 \\ \vdots & \vdots & \vdots \\ 2 & x(N)^2 & x(N)^3 \end{bmatrix}$
---	---	---	---

17. Which of the following formulas computes the covariance for a least squares estimator and a single experiment?  $\hat{\Sigma}_{\hat{\theta}} = \dots$  (all solutions were wrong in the exam, question taken out)

(a) <input type="checkbox"/> $\frac{\  y_N - \Phi_N \hat{\theta} \ _2^2}{N-d} (\Phi_N^T \Phi_N)^{-1}$ corrected solution	(b) <input type="checkbox"/> $\frac{\  y_N - \Phi_N \hat{\theta} \ _2}{N-d} (\Phi_N \Phi_N^T)^{-1}$
(c) <input type="checkbox"/> $\frac{\  y_N - \Phi_N \hat{\theta} \ _2}{N-d} (\Phi_N^T + \Phi_N)$	(d) <input type="checkbox"/> $\Phi_N^+ \sigma_{\epsilon_N}$

18. What is the minimizer  $x^*$  of the convex function  $f : \mathbb{R}_{>0} \rightarrow \mathbb{R}$ ,  $f(x) = -4 \log(x) - \frac{2}{x}$ ? (function not convex, question taken out)

(a) <input type="checkbox"/> $x^* = -2$	(b) <input type="checkbox"/> $x^* = -1/2$	(c) <input type="checkbox"/> $x^* = e^{1/4} - 2$	(d) <input type="checkbox"/> $x^* = 1/2$
---	---	--	--

19. Which of the following is NOT a name of a probability distribution?

(a) <input type="checkbox"/> Uniform	(b) <input type="checkbox"/> Gaussian	(c) <input type="checkbox"/> Newton	(d) <input type="checkbox"/> Laplace
--------------------------------------	---------------------------------------	-------------------------------------	--------------------------------------

20. Given a uniformly distributed random variable  $X$  on the interval  $[-1, 1]$ , regard the following  $X$  - dependent random variables  $Y$ . For one of the examples  $X$  and  $Y$  are uncorrelated. Which one?

(a) <input type="checkbox"/> $y = \sin(x)$	(b) <input type="checkbox"/> $y = \cos(x)$	(c) <input type="checkbox"/> $y = x^3$	(d) <input type="checkbox"/> $y = e^x$
--	--	--	--